



**In Compliance with Oklahoma Statutes, Title 70 §3243**

**Certification of Meningococcal Compliance**

Oklahoma Statutes, Title 70 §3243, requires that all students who are first time enrollees in any public or private postsecondary educational institution in this state and who reside in on-campus student housing shall be vaccinated against meningococcal disease. Institutions of higher education must provide the student or the student's parents or other legal representative detailed information on the risks associated with meningococcal disease and on the availability and effectiveness of any vaccine.

The statute permits the student or, if the student is a minor, the student's parent or other legal representative, to sign a written waiver stating that the student has received and reviewed the information provided on the risks associated with meningococcal disease and on the availability and effectiveness of any vaccine, and has chosen not to be or not to have the student vaccinated.

Student's Name: \_\_\_\_\_ Social Security # \_\_\_\_\_

Birth date: \_\_\_\_\_ Term/Year of enrollment: \_\_\_\_\_

- 1) **I have received and reviewed detailed information on the risks associated with meningococcal disease, and**
- 2) **I have received and reviewed information on the availability and effectiveness of any vaccine (against meningococcal disease), and**
- 3) **I have been vaccinated or   \* I have chosen not to be vaccinated against meningococcal disease. (Please check one) If you have been vaccinated you must provide documentation.**

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

When student is under 18 years of age, the following must also be completed:

**As the parent, guardian or other legal representative, I certify that the student named above is a minor and that I have received and reviewed the information provided and that I have chosen not to have the student vaccinated against meningococcal disease.**

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

\*With this waiver, I seek exemption from this requirement. I voluntarily agree to release, discharge, indemnify and hold harmless Eastern Oklahoma State College, its officers, employees and agents from any and all costs, liabilities, expenses, claims, demands, or causes of action on account of any loss or personal injury that might result from my decision not to be immunized against meningitis. By not being immunized I realize if there is a case of meningitis on campus I will be required to leave campus and remain off campus until it is deemed safe to return.

# Meningococcal Vaccines

## What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

### 1 What is meningococcal disease?

Meningococcal disease is a serious bacterial illness. It is a leading cause of bacterial meningitis in children 2 through 18 years old in the United States. Meningitis is an infection of the covering of the brain and the spinal cord.

Meningococcal disease also causes blood infections.

About 1,000–1,200 people get meningococcal disease each year in the U.S. Even when they are treated with antibiotics, 10–15% of these people die. Of those who live, another 11%–19% lose their arms or legs, have problems with their nervous systems, become deaf, or suffer seizures or strokes.

Anyone can get meningococcal disease. But it is most common in infants less than one year of age and people 16–21 years. Children with certain medical conditions, such as lack of a spleen, have an increased risk of getting meningococcal disease. College freshmen living in dorms are also at increased risk.

Meningococcal infections can be treated with drugs such as penicillin. Still, many people who get the disease die from it, and many others are affected for life. This is why preventing the disease through use of meningococcal vaccine is important for people at highest risk.

### 2 Meningococcal vaccine

There are two kinds of meningococcal vaccine in the U.S.:

- Meningococcal conjugate vaccine (**MCV4**) is the preferred vaccine for people 55 years of age and younger.
- Meningococcal polysaccharide vaccine (**MPSV4**) has been available since the 1970s. It is the only meningococcal vaccine licensed for people older than 55.

Both vaccines can prevent 4 types of meningococcal disease, including 2 of the 3 types most common in the United States and a type that causes epidemics in Africa. There are other types of meningococcal disease; the vaccines do not protect against these.

### 3 Who should get meningococcal vaccine and when?

#### Routine vaccination

Two doses of MCV4 are recommended for adolescents 11 through 18 years of age: the first dose at 11 or 12 years of age, with a booster dose at age 16.

Adolescents in this age group with HIV infection should get three doses: 2 doses 2 months apart at 11 or 12 years, plus a booster at age 16.

If the first dose (or series) is given between 13 and 15 years of age, the booster should be given between 16 and 18. If the first dose (or series) is given after the 16<sup>th</sup> birthday, a booster is not needed.

#### Other people at increased risk

- College freshmen living in dormitories.
- Laboratory personnel who are routinely exposed to meningococcal bacteria.
- U.S. military recruits.
- Anyone traveling to, or living in, a part of the world where meningococcal disease is common, such as parts of Africa.
- Anyone who has a damaged spleen, or whose spleen has been removed.
- Anyone who has persistent complement component deficiency (an immune system disorder).
- People who might have been exposed to meningitis during an outbreak.

Children between 9 and 23 months of age, and anyone else with certain medical conditions need 2 doses for adequate protection. Ask your doctor about the number and timing of doses, and the need for booster doses.

MCV4 is the preferred vaccine for people in these groups who are 9 months through 55 years of age. MPSV4 can be used for adults older than 55.



**4****Some people should not get meningococcal vaccine or should wait.**

- Anyone who has ever had a severe (life-threatening) allergic reaction to a previous dose of MCV4 or MPSV4 vaccine should not get another dose of either vaccine.
- Anyone who has a severe (life threatening) allergy to any vaccine component should not get the vaccine. *Tell your doctor if you have any severe allergies.*
- Anyone who is moderately or severely ill at the time the shot is scheduled should probably wait until they recover. Ask your doctor. People with a mild illness can usually get the vaccine.
- Meningococcal vaccines may be given to pregnant women. MCV4 is a fairly new vaccine and has not been studied in pregnant women as much as MPSV4 has. It should be used only if clearly needed. The manufacturers of MCV4 maintain pregnancy registries for women who are vaccinated while pregnant.

Except for children with sickle cell disease or without a working spleen, meningococcal vaccines may be given at the same time as other vaccines.

**5****What are the risks from meningococcal vaccines?**

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. The risk of meningococcal vaccine causing serious harm, or death, is extremely small.

Brief fainting spells and related symptoms (such as jerking or seizure-like movements) can follow a vaccination. They happen most often with adolescents, and they can result in falls and injuries.

Sitting or lying down for about 15 minutes after getting the shot—especially if you feel faint—can help prevent these injuries.

**Mild problems**

As many as half the people who get meningococcal vaccines have mild side effects, such as redness or pain where the shot was given.

If these problems occur, they usually last for 1 or 2 days. They are more common after MCV4 than after MPSV4.

A small percentage of people who receive the vaccine develop a mild fever.

**Severe problems**

Serious allergic reactions, within a few minutes to a few hours of the shot, are very rare.

**6****What if there is a serious reaction?****What should I look for?**

Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

**What should I do?**

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at [www.vaers.hhs.gov](http://www.vaers.hhs.gov), or by calling **1-800-822-7967**.

*VAERS is only for reporting reactions. They do not give medical advice.*

**7****The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation).

**8****How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines)

---

**Vaccine Information Statement (Interim)  
Meningococcal Vaccine**

10/14/2011

42 U.S.C. § 300aa-26

Office Use Only

